New York State Immunization Requirements for School Entrance/Attendance

NOTES: Children in a prekindergarten setting should be age-appropriately immunized. The number of doses depends on the schedule recommended by the Advisory Committee for Immunization Practices (ACIP).

information for each vaccine. Children who are enrolling in grade-less classes should meet the immunization requirements of the grades for which they are age equivalent. and do not count toward the number of doses listed below. Intervals between doses of vaccine DO NOT need to be reviewed for grades 8 through 12. See footnotes for specific For grades Pre-k through 7, intervals between doses of vaccine should be in accordance with the ACIP-recommended immunization schedule for persons 0 through 18 years of age. (Exception: intervals between doses of polio vaccine need to be reviewed only for grades kindergarten, 1, 6 and 7.) Doses received before the minimum age or intervals are not valid

Dose requirements MUST be read with the footnotes of this schedule.

Pneumococcal Conjugate vaccine (PCV)*	Haemophilus imfluenzae type b conjugate vaccine (Hib)s	Varicella (Chickenpox) vaccine?	Hepatitis B vaccine ^s	Measles, Mumps and Rubella vaccine (MMR)	Polio vaccine (IPV/OPV)*	Tetanus and Diphtheria toxoid-containing vaccine and Pertussis vaccine booster (Tdap) ³	Diphtheria and Tetanus toxoid-containing vaccine and Pertussis vaccine (DTaP/DTP/Tdap) ²	Vaccines
1 to 4 doses	1 to 4 doses	1 dose	3 doses	I dose	3 doses		A 100 CONTROL OF THE PROPERTY	Prekindergarten (Day Care, Head Start, Nursery or Pre-k)
		2 doses	3 do	orage or older	4 doses or 3 doses if the 3rd dose was received at 4 years	Notappicable	5 doses or 4 doses if the 4th do at 4 years of age or o at 4 years of age or o a doses if the series is started or older	Kindergarten through Grade 1
Not applicable	Not applicable	1 dose	3 doses or 2 doses of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 through 15 years of age	2 doses	3 d 88		5 doses or 4 doses if the 4th dose was received at 4 years of age or older or 3 doses if the series is started at 7 years of age or older	Grades 2 through 5
icable	icable	2 doses	ses of adult hepatitis B vaccine (Recombivax ho received the doses at least 4 months apart i the ages of 11 through 15 years of age	or age or ower	4 doses or 3 doses if the 3rd dose was received at 4 years			Grades 6 through 7
		Idose	mbīvax) s apart e		388	1 dos	3 doses	Grades 8 through 12